

Technical Memorandums are used for documentation and timely communication of preliminary results, interim reports, or special-purpose information, and have not received complete formal review, editorial control, or detailed editing.



Species profile of Spanish sardine

(Sardinella aurita)

Allyn G. Johnson

and

Rosalie N. Vaught

August 1986

U.S. DEPARTMENT OF COMMERCE
Malcolm Baldrige, Secretary
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
Anthony J. Calio, Administrator
NATIONAL MARINE FISHERIES SERVICE
William G. Gordon, Assistant Administrator for
Fisheries

I. Introduction

This report is a summary of available information on the biology, fishery, and marketing of Spanish sardines (Sardinella aurita). Emphasis is on the Gulf of Mexico with reference to other areas when particular information may have application to the Gulf of Mexico resource.

The purposes of this report are:

1. To provide a summary of the current knowledge of this species and the resource.
2. To point out the limitations of our current knowledge.

II. Taxonomy and Identification

The Spanish sardine (Sardinella aurita, Valenciennes 1847) is also called round sardinella (English), allache (French) and sardinela atlántica (Spanish). Other names of less common usage are sardina de España, false sardine, sardina, and bang. Previously used scientific synonyms of this species are: Sardinella anchovia, Clupea anchovia, Clupea pseudo-hispanicus, Clupanadon pseudohispanicus, Sardinia anchovia, and Sardinia pseudo-hispanica (Fischer 1978, Hildebrand 1963).

The adult form of this species has the typical sardine appearance (Fig. 1) with the following distinctive characteristics according to Fischer (1978), Robins and Ray (1986):

Body elongate and slender, sometimes cylindrical, but usually a little compressed; belly rounded, with scutes but without a prominent keel. Top of head with 8 to 10 longitudinal fronto-parietal striae on each side; no hypo-maxilla; hind border of gill opening with two fleshy out-growths; gill rakers fine and numerous, increasing with size of fish, anterior rakers on lower limbs of gill arches lying more or less flat. Dorsal fin a little before center point of body; pelvic fins with 9 rays, set below dorsal fin base; anal fin short and well behind dorsal (Fig. 1); last two anal fin rays enlarged, almost like a finlet. Color: back dark blue, changing rather abruptly to silver or brassy on sides; no spots or streaks; fins hyaline. Size: maximum, 29 cm; common to 25 cm.

Spanish sardines can be distinguished from other species in its range by the following characteristics:

Sardinella brasiliensis: anterior rakers on lower limbs of gill arches strongly curled when seen from above in S. brasiliensis (Fig. 2b); narrow bronzy line along upper part of side.

Opisthonema species: last dorsal fin ray filamentous in Opisthonema; also, only 8 pelvic fin rays (9 in Sardinella).

Harengula species: hypo-maxilla present (Fig. 2); also, only 8 pelvic fin rays in Harengula.

Etrumeus teres and Jenkinsia species: no scutes along belly as in Sardinella; also, 14 to 15 branchiostegal rays in Etrumeus (6 in Sardinella).

Lile piquitinga: only 8 pelvic fin rays, belly with sharp keel in L. piquitinga.

Prosvirov and Varea (1969) provided a table of morphometric features of Gulf of Mexico Spanish sardine which they classify as plastic (variable) (Table 1). Hildebrand (1963) using morphometric measurements expressed as percent of standard length from 32 fish (90 to 155 mm total length), also indicated that the morphometric characters of this species are variable (Table 2).

There may be more than one species in the Gulf of Mexico that is referred to as Spanish sardine (Klima 1971). Whitehead (1973) has provided information on several lectotypes, but not from the Gulf of Mexico. Heald and Griffiths (1967), in their paper on age determination of Spanish sardine from Venezuela, refer to two types of Sardinella anchovia from the Gulf of Cariaco. Research is needed to determine the number of species of Spanish sardine, their geographic distribution, and identifying characteristics. This review paper examines the information available on Spanish sardine as if there is only one species, Sardinella aurita.

The larval stages of Spanish sardine have been described by Fahay (1983), (Table 3 and Fig. 3). Jones et al. (1978) provided a series of larval illustrations compiled from various sources (Fig. 4).

III. Distribution

The Spanish sardine occurs throughout the Gulf of Mexico, northward to Massachusetts and southward to Rio de Janeiro, Brazil. This species is also common in the eastern Atlantic, Mediterranean Sea and the western Pacific Ocean (Fischer 1978). In the Gulf of Mexico it is found from along the shore, mainly over sandy bottom, to a depth of 40 meters in the summer, between 40 and 80 meters in spring and fall, and out to a depth of 200 meters in the winter (Fig. 5) (U.S. Dep. Commer. 1985, Houde et al. 1979).

The distribution of larval Spanish sardine for the eastern Gulf of Mexico (west coast of Florida) has been reported by Houde (1973) and Houde et al. (1979). Ichthyoplankton survey catches of 1971-74 (17 cruises) in the eastern Gulf of Mexico were dominated by the families Clupeidae and Gobiidae in water less than 100 m deep. The annual abundance of larvae of Spanish sardine varied by factors greater than 20 times between years in the area studied (Figs. 6-8). Since clupeid eggs hatch 20 to 45 hours after spawning at water temperatures of the eastern

Gulf of Mexico (Houde 1973), we assume the larvae originated near the area of capture. The greatest abundance occurred in the summer and fall (May-October) (Fig. 7).

Very little information is available on larval distribution of Spanish sardine in the western Gulf of Mexico. A few larvae have been collected in waters as deep as 180 m in surface ichthyoplankton sampling in March through December (Finucane et al. 1977, 1978).

The adult Spanish sardine is "normally found close to the coast, well within the 50-fathom curve, but it has been caught as far offshore as the 200-fathom line in the Atlantic" (Klima 1971). Low (1973) reported that Spanish sardine (40 to 160 mm total length) occurred along with *H. pensacolata* and *O. oglinum* over grassbeds in Biscayne Bay, Florida (east coast) from May into the fall, thus indicating this species will occur in shallow waters. The distribution is generally along the shoreline to water 100 fathoms (183 m) deep, (Figs. 5 and 9). The fish are generally in the upper 40 fathoms (74 m) of the water column (Tables 4 and 5). The species has been caught in waters 13° to 31°C surface temperature and 17° to 28°C bottom temperature (Tables 6 and 7). Commercial concentrations of sardines have been reported 90 miles from shore at depths of 40 to 80 m off western Florida (Tampa area) by Prosvirov and Varea (1969). Another area of high concentration that has been reported is Campeche Bank where 4.7 to 5.3 tons/hr (using bottom trawl) have been caught (Sal'nikov 1965).

The species appears to form schools which migrate in the surface layers along the continental shelf. This species makes diurnal vertical migrations (keep to the bottom during day and rise to midwater at dusk). This upward movement appears to depend on illumination (Prosvirov 1967). Seasonality of movements is not well defined. Hildebrand (1963) believed the fish made a southward migration along the Atlantic coast in the fall. Low (1973) reported fish in Biscayne Bay, Florida in fall and winter but not spring and summer.

IV. Biology

Reproduction

The reproduction of Spanish sardine has not been examined thoroughly. The following excerpt from Reintjes (1980) summarizes the current information on Spanish sardine.

"Spawning occurs in the open waters of the Gulf, although the exact areas of spawning are difficult to determine. According to Houde et al. (1976, 1979), most spawning in the eastern Gulf occurs over wide areas where depths are between 10 and 50 m, although some eggs and larvae have been recovered where depths were 200 m. In the western Gulf, larvae have been reported from as far out as the Continental Shelf off Texas (Finucane, Collins, and Barger 1978). Eggs and larvae have been collected in the eastern Gulf in all

seasons, but are not abundant from May to September (Houde et al. 1976, 1979), and in the western Gulf from April to October (Finucane, Collins, and Barger 1978). Since spawning occurs over a long period, a given year class will have a wide range in lengths (Prosvirov and Varea 1969). Elsewhere, off the northeast coast of Venezuela, Lopez (1972) reported that eggs were most abundant from August to November and in February and May, while Simpson and Griffiths (1967) stated that spawning occurred year round but was most intense during the upwelling season from December to April.

Only a few larvae have been described. Houde and Fore (1973) show figures of a larvae 7.8 mm and 19.3 mm, while Simpson and Gonzalez (1967) show detailed illustrations of developing embryos but not larvae. Larvae from off Brazil that may have been Spanish sardine were described by Aboussouan (1969); larvae of S. aurita were described by D'Ancona (1956).

Fecundity estimates for Spanish sardine in the Gulf are sparse. For eight females, 14.4 to 15.5 cm SL, collected near Palm Beach, Florida in 1971, estimates ranged from 30,718 to 86,269 (Martinez 1972). In the Mediterranean fecundity ranged from 3,710 to 48,600 depending on size of females (Ben-Tuvia 1960); off West Africa it ranged from 10,000 to 270,000."

Martinez (1972) provided more detailed information on fecundity of Spanish sardine from the west coast of Florida (Biscayne Bay), which is summarized in Tables 8 and 9 and Figure 10. The most recent work on reproduction of Spanish sardine in Florida waters was by Grall (1984). She reported on gonad index, fecundity and maturation (summarized in Tables 10 to 13, Figures 11 to 15). She found that although sex ratios varied considerably from month to month, the overall sex ratio in the Gulf of Mexico in 1982 was 1:1. No specific information was found on factors influencing spawning or spawning behavior.

Population Parameters

The most recent data on population parameters were reported by Grall (1984) from fish of northwest Florida and southeast Florida.

The length frequency of Spanish sardine from commercial fisheries for the above areas are presented in Figures 16 and 17. Mean fork length and weight by month showing smaller and lighter fish being caught in the fall are presented in Figure 18 and Tables 14 and 15. The length-weight relationship is described in Figure 19 and Tables 16 and 17.

Age determination was made on Spanish sardine using marks (deposited October - April) on otoliths. This information and growth parameters are presented in Tables 18 to 22 and Figure 20. Comparisons of growth parameters are presented in Table 23.

Mortality and survival estimates using the above data were developed and shown along with yield-per-recruit estimates in Tables 24 and 25 and Figure 21.

Additional age and growth information on Spanish sardine has been developed by Scherbich (1981) for the southeast Atlantic and Heald and Griffiths (1967) for Venezuela using scales as the age determination structure. Scherbich reported ages up to 6 yrs old and Heald and Griffiths up to 5 yrs old.

Houde et al. (1979) provided information on larval Spanish sardine abundance and mortality. The annual abundance varied considerably from year to year off the west coast of Florida (Fig. 22). His mortality coefficient for larval Spanish sardine was 0.3909.

Ecological Associations

Little is known concerning the food habits of Spanish sardine. They apparently feed primarily on planktonic copepods (Hildebrand 1963, Böhlke and Chaplin 1968, Low 1973).

Predators of Spanish sardine are numerous. Manooch (1984) reported that little tunny (*Euthynnus alletteratus*), gag (*Mycteroperca microlepis*), and king mackerel (*Scomberomorus cavalla*) eat Spanish sardine. Other references to predators eating this species are provided in the papers by Saloman and Naughton, which are summarized in Table 26.

Behavior and Environmental Relationships

The behavior and environmental relationships of Spanish sardine have not been investigated in detail. Klima and Wickham (1971) reported that Spanish sardine were attracted to artificial structures. Maksimov (1976) reported that defensive behavior is taken by this species in response to trawls. This behavior is usually expressed in the "zone of critical narrowing" of the trawl and consists of fright, optomotor reaction, etc., which causes the fish to leave the trawl. Klima (1972) reported laboratory observations on Spanish sardine to different voltage and direct current pulse rates (1.5 to 3.0 v/10 cm at 15 to 45 pulses/sec with a duration of 1.0 m sec). Zei (1969) reported that this species could be attracted by artificial light at night and that high catches were associated with low temperature in the Gulf of Guinea. He also reported that successful purse seine catches were dependent on the fish being assembled in catchable numbers which was greater in periods of upwelling than in periods of thermal stability. Aleem (1972) reported decreases in catches of Spanish sardine because of construction of the Aswan high dam, which caused a drop in nutrients from the Nile River, resulting in the disappearance of phytoplankton blooms. This may indicate positive response on the part of Spanish sardine to food concentrations. Low (1973) also associated the appearance of this species with food concentrations. A complex environmental (temperature and predator-prey) interaction may exist off the west coast of Africa

between Spanish sardine and gray triggerfish (Balistes capriscus) where cold upwellings and high salinity favor the sardine and the opposite favor the triggerfish which feeds on benthos and pelagic organisms, including clupeid eggs and larvae (FAO 1980).

Stocks

The identification of stocks has not been adequately examined for Spanish sardine in the Gulf of Mexico area. Prosvirov (1967) suggested that Spanish sardine in the Gulf of Mexico has a number of local populations. Johnson (unpublished ms-a) reported that 8.6% of 35 enzymatic loci (starch gel electrophoresis) were polymorphic and estimated that the maximum proportion of the genome heterozygous was 0.051. This indicates that protein polymorphism may be useful for stock separation. Other reports on electrophoretic polymorphism in serum proteins (from Guinea and Mauritania) suggest they may be useful to separate stocks (Baron 1972, 1973a, 1973b). Preliminary work on morphometrics by Johnson (unpublished ms-b) showed that stepwise discriminant analysis of multiple measurements could correctly classify fish 97.3% of the time between three collection areas (Panama City, Cortez, and Fort Pierce) in Florida, which indicated that more than one stock may occur in the southeastern U.S. waters. Latini and Pettorossi (1977) used meristics to separate various collections of Mediterranean Spanish sardines. Meristics have not been examined for fish from the Gulf of Mexico, but may be useful to separate stocks of fish.

V. Fisheries and Catching Methods

The commercial fishery for Spanish sardine can be briefly summarized by the following excerpt from Reintjes (1980).

"The largest fishery in the western Atlantic is off Venezuela, where annual landings of more than 40,000 tons have been reported. Mexico has reported landings of 1,000 tons in some years, and Sal'nikov (1965) and Sokolova (1965) have reported Spanish sardine in catches from Campeche Banks. An estimated 1,000 tons are landed annually in a small seine fishery from May to October in the Florida panhandle from Port St. Joe to Pensacola. Most of this catch is used for bait."

The Spanish sardine supports the most important commercial fishery in Venezuela where the major gear type is the beach seine. Landings have averaged 40,000 mt annually from 1963 to 1967 having grown from 4,600 mt (Simpson 1971). Off Jamaica, Spanish sardines are caught by a variety of gear types (Fig. 23 and 24) - gill net, beach seine, lift nets, and cast nets (Harvey 1982).

In the northeastern Gulf of Mexico the Spanish sardine is caught by the coastal pelagic purse seine. The purse seine vessels average 45 to 65 ft. in length. They fish by means of a large net that encircles the school. The bottom of the net can be "pursed", thereby completely

enclosing the fish. The net is slowly hauled in, and the fish are brought aboard by using a large brail or dip net. A detailed description of this operation can be found in Roithmayr 1983 (see Fig. 25). This source also gives the fishing grounds for the purse seine fleet as: from the Mississippi River to Apalachicola, Florida; also, west-central Florida from St. Petersburg, to Sarasota (Fig. 26). Principal ports for the purse seiners include Yscloskey, La.; Pascagoula, Miss.; and in Florida: Destin, Panama City, Port St. Joe, and Cortez. A drawback of this fishing method is some loss of fish quality, due to the abrasion that takes place while the fish are confined in the net.

Spanish sardine comprise a minor portion of the catch of this gear type (for example - Table 27); however, the amount of this species in a catch depends on the the vessel's target, based on market demands.

Additional information on exploratory purse seining is available in a report by Harrell (1982). She provides the cost of converting a 78 ft steel shrimp trawler (150,000 lb hold capacity) to a dual purpose drum seine/pair trawling vessel (Table 28). The result of fishing tests demonstrated that drum seining must produce 30,000 pounds of fish per day in order to be profitable (catch valued at 15 to 25 cents per pound) and can give a 21% return on investment (Tables 28, 29, and 30).

Centaur Associates (1981) provided the following excerpt which described the northern Gulf of Mexico fishery.

"Most of the coastal herring catch is currently taken from inshore waters. Two types of gear are utilized in the directed fishery. Haul or beach seines are fished with boats ranging in size from 23 to 50 feet in length. The seines vary in length from 400 to 750 yards. Purse seines are the second gear type used. These are fished with vessels ranging from 45 to 65 feet in length. The nets are approximately 750 yards long and generally 45 to 60 feet deep. Approximately 6 purse seine boats were reported fishing these species in the Gulf of Mexico. Purse seine boats land greater quantities than haul seines.

A total of 30 to 35 boats and vessels currently participate in the directed fishery for coastal herrings. According to Reintjes (1979), most of these are located on the west coast of Florida. Alabama is the primary landing area for coastal herrings in the northern Gulf. Thread herring and blue runners are reported in the Alabama commercial catch. However, most, if not all, of the landings of these species in the northern Gulf are catches incidental to other fishing activities, primarily the menhaden, shrimp and ground fish fleets....The resource potential for a directed coastal herring fishery in the northern Gulf appears good, but current processing and demand problems will have to be overcome. Development of a directed fishery in the northern Gulf would likely place a greater demand on port facilities for relatively small boats and vessels if the type presently used in Florida are fished in the

northern Gulf. Development of this fishery, however, would not result in a proportional increase in number of boats and vessels since many of the existing craft could expand into the fishery. many of the Florida boats and vessels who fish for bait also fish for other existing foodfish species such as pompano, mullet and the mackerel."

An example of the value of the purse-seine-caught fish was provided by Uebelhoer (1984), where Spanish sardine made up 11% of the return for 26 sets (purse seine catches) (Table 31).

Whether or not the abundance of, and markets for, Spanish sardine can support full development of this type of fishery is unknown, but Houde (1976) suggested that there may be about 250,000 metric tons of Sardinella sp. in the eastern Gulf of Mexico. Bullis and Carpenter (1968) suggested that 8,448 million pounds are available for harvest in the coastal pelagic fish resource (herring group) (Table 32). The coastal herrings can be broken down further into their respective components which are presented in Table 33 (Reintjes 1979).

VI. Composition and Marketing

Hale (1984) found the proximate composition of Spanish sardine from the gulf to be: protein, 20.90%; fat, 2.42%; moisture, 74.73%; ash, 2.69%.

The quality of a fish species with regard to oil, protein, usable meat, storage etc., greatly influences the desirability as a resource. Various sources have made studies on the quality of Spanish sardine (Table 34). The fatty acid and lipid classes and changes due to canning was studied by Hale and Brown (1983). They reported no significant changes in these traits as a result of canning; however, the vegetable oil used in packing had major effects on the fatty acid profile (increases in lenoleic acid). The chemical composition between raw and canned fish showed an increase in percent protein and lipid and a decrease in percent moisture in the canned product compared to the raw fish (Tables 35, 36, 37 and 38).

Another study showed that fish held in refrigerated sea water (brine spray system) were of higher quality than those held in chilled sea water (slush system) or ice during short-term storage. However, excessive saltiness and rancidness developed by the third day in refrigerated sea water and by the fourth day in chilled sea water and ice. Whole and headed and gutted fish stored frozen showed little deterioration for up to three months. Of the three holding methods, only chilled sea water produced an acceptable canned product for fish that were held for more than one or two days. Frozen fish could be made into good canned products even after two months of frozen storage. In order to produce a quality product, careful handling was required. Off loading by brailing was preferred over pumping (ABIC International Consultants, Inc. 1984).

Bushardt and Williams (1983) have examined the problems involved in the packaging of underutilized species for marketing. They stress the need for development of high quality, identifiable boxes and an industrial standard. They suggest that Spanish sardine be packed in 3-mil poly bags in banded 23" x 13-1/2" x 5-7/8" single wall, 200 lb, corrugated, water resistant cartons.

The current landings of Spanish sardine of 1,265 mt for the United States and 56,082 mt for the total area (FAO 1984) indicate that considerable amounts remain for harvesting. Combs (1978) indicated that in the United States, current use of this species is as bait and frozen animal food, but that Spanish sardine has potential for use as meal and as canned products and also as a substitute for menhaden and other sardine products. The constraints on this fishery have been the wide annual fluctuations in resource availability and the fact that the fish spoils quickly.

Perkins (1981) provided an economic assessment of the potential sardine fishery for the northern Gulf of Mexico. He reports very positive reaction to and high preferences for canned Gulf of Mexico Spanish sardines. Commercial fishermen can profitably produce large volumes of fish for a price of \$160 to \$200 per ton which would yield an ex-vessel value of 10.8 million dollars. This volume would yield a retail canned value of 37.7 million dollars and would generate an economic impact of 75 to 113 million dollars annually. His report also gives a detailed description of an experimental canning operation for Spanish sardine at an established seafood canning plant in Mississippi, using various packing mediums. Regarding demand, he states that the U.S. consumption of sardines is low, averaging only 0.4 pounds per capita per year over a seven year period (Table 39). However, it is expected that consumption would rise with some reduction in price (Figs. 27 and 28; Tables 40 and 41).

The number of firms exporting underutilized species from the Gulf of Mexico and south Atlantic area have been examined by Prochaska and Cato (1981) and Centaur Associates (1981). Their information is summarized in Tables 42 and 43.

The following are two personal communications about the market prospects of Spanish sardines from marketing specialists.

From M.E. Smith (National Marine Fisheries Service, Pascagoula, MS, April 16, 1985):

Domestic Market - Other than as bait for recreational fishermen and to lesser extent for zoos, Mr. Smith is not aware of any other use for the fish, fresh or frozen. From the standpoint of eye appeal and edibility, Spanish sardines from the Gulf of Mexico should be well received as a canned item. The same species is canned and imported from Venezuela, and has been seen on shelves in markets at prices as low as 40 cents per 4-ounce can. From the standpoint of size, the gulf species

should make a pack having as much or more eye appeal than Atlantic herring. With the traditional 4-ounce flat can of domestic sardines retailing at the competitive price of 55 cents to 58 cents, it would be more realistic to can the product in an existing facility rather than construct one that would be new. There are several shrimp canneries along the gulf coast that are in dire need of expanding their productivity. The only deterrent to their including sardines would be if the production season for Spanish sardines overlapped or interfered with the canning of small or medium shrimp. A shortage of the resource in New England would probably stimulate an investigation of the Gulf of Mexico stocks.

Export Market - Canned sardines have worldwide acceptance, and this is best seen by the number of inquiries that are received, especially from southeast Asia, about the availability of the product. In this area Japan is a major exporter of canned Spanish sardines, but the product is normally not canned until there is an oversupply, at which time the price on fresh sardines dips to 4 cents per pound. With such a low cost raw material, it is felt that U.S. canneries could not be competitive; this suggests that potential buyers are merely shopping for lower prices. Normally, fresh sardines in southeast Asia are used by low income groups and are also seen dried for the same purpose, wholesaling at approximately 30 cents per pound.

From Richard Raulerson (Southeast Regional Office, NMFS, St. Petersburg, FL, May 25, 1985): Spanish sardine has great potential for fishery development and has been canned for human food in the U.S. for 100 yrs. Currently the fish is used mostly as bait, but has greater potential value as a canned fish. There is a problem in canned fish with an unattractive color in the oil, but this could be worked out technologically. There is a substantial fishery in Cortez, Florida; the fishing industry there also uses it as a home canning product. There once was a canning plant in Pascagoula, Mississippi, run by a Brazilian firm. Major problems for development are competition with the better known canned sardine products, such as the Norwegian product, and trade and tariff restrictions. Spanish sardine is highly used as a canned product in South America.

Spanish sardine also shows potential for use as surimi or a minced fish product. Feasibility research showed that processing for minced fish provided a net yield of 42.6%; processing for surimi provided a net yield of 25.3%. Although the surimi provided a lower net yield, it was reported to have better flavor, color and texture than the minced fish product (Thrash 1985).

VII. Information Needs

The more important areas in which information is needed are:

1. Species identification and geographic distribution. Are there more than one species being called Spanish sardine in the Gulf of Mexico?

If so, what are the identification characteristics and ranges of the species?

2. Accurate estimates of abundance and location of resource, especially in the western Gulf of Mexico.

3. Stock identification, separation, ranges and biological parameters of each stock. Estimates of other resource user needs, so adequate resource allocation can be made. For example, if we take x amount of Spanish sardine, what effect does this removal have on king mackerel and other predators of Spanish sardine?

4. Fishing methods and how to locate this species accurately and effectively catch it. We need to apply innovative and new technology to this problem. For example, Harrell (1982) reported using night flying aircraft to spot groups of fish by the bioluminescent glow caused in the water around the fish.

5. Development of improved methods of handling this species to maintain a quality product. This includes improvements in ship board storage between catching and unloading, and standardization of an industrial "pack" (i.e., development of a universally accepted box for frozen storage of this product).

6. Test marketing of canned and other forms of this species to obtain an accurate evaluation of the consumer response to Spanish sardine. Additional information needs on this species were expressed in Christmas et al. (1985), which are summarized in (Figure 29).

LITERATURE CITED

- ABIC INTERNATIONAL CONSULTANTS, INC. 1984. Effect of preprocess handling and holding conditions on the quality of fresh, frozen and canned Spanish sardines. Prep. for Gulf & So. Atl. Fish. Devel. Found. Inc., NMFS Award #22-09-30875. ABIC Int. Consult., Pine Brook, NJ, pag. var.
- ABOUSSOUAN, A. 1969. Sur une petite collection de larves de Teleosteens recoitee au large du Brasil (Campagne "Calypso" 1962). Vie Milieu 20:595-610.
- ALEEM, A.A. 1972. Effect of river outflow management on marine life. Mar. Biol. 15:200-208.
- BARON, J.C. 1972. Preliminary studies on the blood of Sardinella from the west African coast. XIIth Europ. Conf. Anim. Blood Groups Biochem. Polymorph., Bp., pp. 593-595.

- BARON, J.C. 1973a. Les esterases du serum de Sardinella aurita Valenciennes, 1847, in C.V. application a l'etude des populations. Cah. O.R.S.T.O.M., Ser. Oceanogr. 11(4):389-418.
- BARON, J.C. 1973b. Note sur les proteines seriques de Sardinella aurita Val., 1847 in C.V. Cah. O.R.S.T.O.M., Ser. Oceanogr. 11(2):133-169.
- BEN-TUVIA, A. 1960. Synopsis of biological data on Sardinella aurita of the Mediterranean Sea and other waters. FAO Fish. Biol. Synop. 14:287-312.
- BOELY, T. 1982. Etude du cycle sexuel de la sardinelle ronde (Sardinella aurita Val. 1847) au Senegal. Oceanogr. Trop. 17(1):3-13.
- BÖHLKE, J.E. and C.C.G. CHAPLIN. 1968. Fishes of the Bahamas and adjacent tropical waters. Livingston Pub. Co., Wynnewood, Pa., 771 p.
- BULLIS, H., Jr. and J.S. CARPENTER. 1968. Latent fishery resources of the central west Atlantic region. In: D.W. Gilbert (ed.). The future of the fishing industry in the United States. Univ. Wash. Pub. Fish., new ser., IV, Seattle, WA., 346 p.
- BUSHARDT, S.C. and D.C. WILLIAMS, Jr. 1983. The packaging of underutilized species of fin fish for export from the Gulf and South Atlantic region: problems, constraints, and recommendations. Prep. for Gulf & So. Atl. Fish. Devel. Found., Inc., NMFS Award #NA82-GA-H-00007, 42 p.
- CENTAUR ASSOCIATES. 1981. Port and harbor infrastructure assessment relating to the development of underutilized finfish of the southeast; Alabama report. Prep. for the Gulf & So. Atl. Fish. Devel. Found. Inc., Tampa, FL, 161 p.
- CHRISTMAS, J.Y., D.J. ETZOLD, T.D. McILWAIN and L.B. SIMPSON. 1985. Marine fisheries initiative: Gulf of Mexico phase. Gulf States Marine Fisheries Commission, Ocean Springs, MS, var. pag.
- COMBS, E.R. 1978. A study to determine the export and domestic markets for currently underutilized fish and shellfish. U.S. Dep. Commer. contract no. MO-A01-78-00-4037.
- D'ANCONA, U. 1956. Sottordine Clupeoidei, p. 1-16. In: Fauna e Flora del Golfo de Napoli; Uova, larve e stadi giovanili di Teleostei. Stazione Zoologica de Napoli, Mono. 38.

- EL-MAGHRABY, A.M., G.A. BOTROS and I.A.M. SOLIMAN. 1970. Age and growth studies on Sardinella maderensis Lowe and Sardinella aurita Cuv. and Val. from the Mediterranean Sea at Alexandria (U.A.R.). Bull. Inst. Oceanogr. Fish., Cairo 1:49-82.
- FAHAY, M. P. 1983. Guide to the early stages of marine fishes occurring in the western North Atlantic Ocean, Cape Hatteras to the southern Scotian Shelf. J. Northw. Atl. Fish. Sci. 4:1-423.
- FAO. 1984. Yearbook of fishery statistics, v. 54, 1982. FAO, Rome, 393 p.
- FAO FISHERY COMMITTEE FOR THE EASTERN CENTRAL ATLANTIC. 1980. Report of the ad hoc working group on Sardinella off the coast of Ivory Coast-Ghana-Togo, 2-7 July 1979, Tema, Ghana, and 24-28 Sept. 1979, Abidjan, Ivory Coast. FAO, Rome, CECAP/ECAF Ser. 80/21, 72 p.
- FINUCANE, J.H., L.A. COLLINS and L.E. BARGER. 1978. Determine the effects of discharges on seasonal abundance, distribution and composition of ichthyoplankton in the oil field. 157 p. In: W.B. Jackson (ed.), Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1977-1978. NOAA Rept. to EPA IAG-D5-E696-EO, NMFS, Southeast Fisheries Center, Galveston, TX.
- FINUCANE, J.H., L.A. COLLINS, L.E. BARGER and J.D. McEACHRAN. 1978. Ichthyoplankton/mackerel eggs and larvae. NOAA final report to BLM. Environmental studies of the south Texas outer continental shelf, 1977. SEFC, Galveston Laboratory, TX, 504 p.
- FINUCANE, J.H., L.A. COLLINS and J.D. McEACHRAN. 1977. Ichthyoplankton/mackerel eggs and larvae. NOAA final report to BLM. Environmental studies of the south Texas outer continental shelf, 1976. SEFC, Galveston Laboratory, TX, 484 p.
- FISCHER, W. (ed.). 1978. FAO species identification sheets for fishery purposes. Western central Atlantic (Fishing area 31). v. II. Rome, FAO, var. pag.
- GHENO, Y. 1975. Nouvelle etude sur la determination de l'age et de la croissance de Sardinella aurita Val. dans la region de Pointe Noire. Cah. O.R.S.T.O.M. Ser. Oceanogr. 13(3):251-262.
- GRALL, C. 1984. A study of the biology of the Spanish sardine, Sardinella aurita, in Florida waters. M.S. Thesis, Univ. of Miami, 107 p.
- HALE, M.B. 1984. Proximate chemical composition and fatty acids of three small coastal pelagic species. Mar. Fish. Rev. 46(1):19-21.

- HALE, M.B. and T. BROWN. 1983. Fatty acids and lipid classes of three underutilized species and changes due to canning. Mar. Fish. Rev. 45(4-6):45-48.
- HARRELL, M.R. 1982. Results of exploratory drum seining efforts. Gulf and South Atlantic Fisheries Development Foundation, Inc. NMFS (NOAA) cooperative agreement NA80-GA-H-0005, var. pag.
- HARVEY, G.C. McN. 1982. Artisanal fisheries for herrings in Jamaica. Proc. Gulf Caribb. Fish. Inst. 34:141-148.
- HEALD, E.J. 1966. Age determination of the Spanish sardine (Sardinella anchovia) from the Gulf of Cariaco, Venezuela. M.S. Thesis, Univ. Miami, 98 p.
- HEALD, E.J. and R.C. GRIFFITHS. 1967. The age determination, from scale readings, of the sardine, Sardinella anchovia, of the Gulf of Cariaco, eastern Venezuela. Venezuela Minist. Agric. Cria, Invest. Pesq., Ser. Rec. Explot. Pesq. 1(10):374-446.
- HERZBERG, A. and R. PASTEUR. 1969. Proximate composition of commercial fishes from the Mediterranean Sea and the Red Sea. Fish. Ind. Res. 5(2):39-65.
- HILDEBRAND, S.F. 1963. Genus Sardinella Cuvier & Valenciennes 1847. In: H.B. Bigelow (ed.) Fishes of the western North Atlantic. Sears Found. Mar. Res., Yale Univ., Mem. I, pt. III:397-411.
- HOUDE, E.D. 1973. Estimating abundance of sardine-like fishes from egg and larval surveys, eastern Gulf of Mexico: preliminary report. Proc. Gulf Caribb. Fish. Inst. 25:68-78.
- HOUDE, E.D. 1976. Abundance and potential for fisheries development of some sardine-like fishes in the Gulf of Mexico. Proc. Gulf Caribb. Fish. Inst. 28:73-82.
- HOUDE, E.D., S.A. BERKELEY, J.J. KLINOVSKY, and C.E. DOWD. 1976. Ichthyoplankton survey data report. Univ. of Miami Sea Grant Program, Sea Grant Tech. Bull. 32, 193 p.
- HOUDE, E.D. and P.L. FORE. 1973. Guide to identity of eggs and larvae of some Gulf of Mexico clupeid fishes. Fla. Dept. Nat. Resour., Mar. Res. Lab. Leaflet. Ser. 4(23), 14 p.
- HOUDE, E.D., J.C. LEAK, C.E. DOWD, S.A. BERKELEY and W.J. RICHARDS. 1979. Ichthyoplankton abundance and diversity in the eastern Gulf of Mexico. A report to the Bureau of Land Management, contract #AA550-CT7-28. Univ. of Miami, 546 p.
- JOHNSON, A.G. Unpubl. ms(a). Electrophoretic patterns of proteins in Spanish sardine (Sardinella aurita).

- JOHNSON, A.G. Unpubl. ms(b). Report on morphometric analysis of Spanish sardine.
- JONES, P.W., F.D. MARTIN and J.E. HARDY, Jr. 1978. Development of fishes of the Mid-Atlantic Bight, vol. I: Acipenseridae through Ictaluridae. U.S. Fish Wildl. Serv., Biol. Serv. Prog. FWS/OBS-78/12, 366 p.
- KLIMA, E.F. 1971. Distribution of some coastal pelagic fishes in the western Atlantic. Commer. Fish. Rev. 33(6):21-34.
- KLIMA, E.F. 1972. Voltage and pulse rates for inducing electrotaxis in twelve coastal pelagic and bottomfishes. J. Fish. Res. Board Can. 29(11):1605-1614.
- KLIMA, E.F. and D.A. WICKHAM. 1971. Attraction of coastal pelagic fishes with artificial structures. Trans. Amer. Fish. Soc. 100(1):86-99.
- KRZEPTOWSKI, M. 1981. Growth characteristics of gilt sardine (Sardinella aurita Val., 1847) from Senegal. Acta Ichthyol. Piscatoria 11(2):27-38.
- LATINI, E. and L. PETTOROSSO. 1977. Osservazioni sui caratteri biometrici e meristici di Sardinella aurita Valenciennes nell'alto Adriatico (Pisces, Clupeidae). Quad. Lab. Tecnol. Pesca 2(3):143-151.
- LOPEZ, R.H. 1972. Distribucion y abundancia estimada de huevos de la sardina (Sardinella anchovia) en la region oriental de Venezuela, 1968-1969. Minist. Agric. Cria Venezuela, Proy. Inv. Desarr. Pesq. MAC-PNUD-FAO Inform. Tecnico 46, 12 p.
- LOW, R.A., Jr. 1973. Shoreline grassbed fishes in Biscayne Bay, Florida, with notes on the availability of clupeid fishes. M.S. Thesis, Univ. Miami, 145 p.
- MAKSIMOV, Y.M. 1976. Characteristic features of the defensive behavior of pelagic fish from the bottom trawl zone of the Gulf of Mexico. J. Ichthyol. 16(3):478-484.
- MANOOCH, C.S., III. 1984. Fisherman's guide--Fishes of the southeastern United States. N.C. St. Mus. Nat. Hist., Raleigh, N.C., 362 p.
- MARTINEZ, S. 1972. Fecundity, sexual maturation and spawning of scaled sardine (Harengula pensacolae). M.S. Thesis, Univ. Miami, 51 p.
- MATSUURA, Y. 1977. O ciclo de vida da sardinha-verdadeira (introducao a oceanografia pesqueira). Pub. Esp. Inst. Oceanogr., Sao Paulo 4:1-146.

- NAUGHTON, S.P. and C.H. SALOMAN. 1984. Food of bluefish (Pomatomus saltratrix) from the U.S. south Atlantic and Gulf of Mexico. NOAA Tech. Memo. NMFS-SEFC-150, 37 p.
- PAULY, D. 1980. On the interrelationships between natural mortality, growth parameters and mean environmental temperature in 175 fish stocks. J. Cons. Int. Explor. Mer 39(2):175-192.
- PERKINS, G.B. 1981. Economic assessment of a potential sardine fishery in the northern Gulf of Mexico. Miss. St. Univ., Food and Fiber Center, 22 p.
- PROCHASKA, F.J. and J.C. CATO. 1981. Developing export markets for Gulf of Mexico and south Atlantic seafood products. In: R. Nickelson, II (comp.). Proceedings of the sixth annual tropical and subtropical fisheries technological conference of the Americas. Texas A&M Univ., Sea Grant Pub. TAMU-SG-82-101:47-59.
- PROSVIROV, E.S. 1967. Distribution of Sardinella anchovia Cuvier and Valenciennes in the Gulf of Mexico and some aspects of its biology (in Russian). p. 185-196. In: A.S. Bogdanov (ed.). Sovetsko-Kubinskije rybokhozyaistvenne issledovaniya, VNRIO-CIP, Vypusk II. Izdatel'stvo "Pischevaya Promyshlennost", Moskva.
- PROSVIROV, E.S. and A. VAREA. 1969. Sardinella anchovia Valenciennes of the Gulf of Mexico. p. 274-281. In: A.S. Bogdanov (ed.). Soviet-Cuban fishery research. Israel Prog. for Sci. Trans., Jerusalem, 350 p.
- REINTJES, J.W. 1979. Coastal herrings and associated species—a profile of species or groups of species, their biology, ecology, current exploitation with economic and social information. Prep. for the Gulf and So. Atl. Fish. Manage. Counc. Southeast Fisheries Center, Beaufort Lab., Beaufort, NC, 170 p.
- REINTJES, J.W. 1980. Marine herring and sardine resources of the northern Gulf of Mexico. In: M. Flandorfer and L. Skupien (eds.). Proceedings of a workshop for potential fishery resources of the northern Gulf of Mexico. Miss.-Ala. Sea Grant Consort. pub. no. MASGP-80-012:4-17.
- ROBINS, C.R. and G.C. RAY. 1986. A field guide to Atlantic coast fishes of North America. Houghton Mifflin Co., Boston, MA, 354 p.
- ROITHMAYR, C.M. 1983. Purse seine fishing for coastal pelagic species in the northern Gulf of Mexico. In: R. Nickelson, II (ed.). Proceedings of the eighth annual tropical and subtropical fisheries conference of the Americas. Texas A&M Univ. Sea Grant Publ. TAMU-SG-83-112:85-101.

- SAL'NIKOV, N.E. 1965. Fishery research in the Gulf of Mexico and the Caribbean Sea, p. 78-171. In: A.S. Bogdanov (ed.). Sovetsko-Kubinskie rybokhozyaistvennye issledovaniya, VNIRO-CIP, Vypusk I. Izdatel'stvo "Pishchevaya Promyshlennost", Moskva. Soviet-Cuban Fishery Research, Israel Program for Scientific Translations, Jerusalem, 1969.
- SALOMAN, C.H. and S.P. NAUGHTON. 1983. Food of king mackerel, Scomberomorus cavalla, from the southeastern United States including the Gulf of Mexico. NOAA Tech. Memo. NMFS-SEFC-126, 25 p.
- SALOMAN, C.H. and S.P. NAUGHTON. 1983. Food of Spanish mackerel, Scomberomorus maculatus, from the Gulf of Mexico and southeastern seaboard of the United States. NOAA Tech. Memo. NMFS-SEFC-128, 22 p.
- SALOMAN, C.H. and S.P. NAUGHTON. 1984. Food of crevalle jack, (Caranx hippos) from Florida, Louisiana, and Texas. NOAA Tech. Memo. NMFS-SEFC-134, 34 p.
- SCHERBICH, L.V. 1981. Age determination methods, growth equations, and age and length composition for Sardinella aurita (Walbaum 1847) in the southeast Atlantic. Collect. Sci. Pap. Intl. Comm. SE. Atl. Fish. (ICSEAF 1981) pt. II:245-251.
- SIDWELL, V.D., P.R. FONCANNON, N.S. MOORE and J.C. BONNET. 1974. Composition of the edible portion of raw (fresh or frozen) crustaceans, finfish, and mollusks. I. Protein, fat, moisture, ash, carbohydrate, energy value, and cholesterol. Mar. Fish. Rev. 36(3):21-35.
- SIDWELL, V.D. 1981. Chemical and nutritional composition of finfishes, whales, crustaceans, mollusks, and their products. NOAA Tech. Memo. NMFS F/SEC-11, 432 p.
- SIMPSON, J.G. 1971. The present status of exploitation and investigation of the clupeoid resources of Venezuela. FAO Fish. Rep. 71.2:263-278.
- SIMPSON, J.G. and G.G. GONZALEZ. 1967. Some aspects of the early life history and environment of the sardine Sardinella anchovia, in eastern Venezuela. Venezuela Minist. Agric. Cria, Invest. Pesq. Ser. Rec. Explot. Pesq. 1(2):38-93.
- SIMPSON, J.G. and R.C. GRIFFITHS. 1967. The fishery resources of Venezuela and their exploitation. Venezuela Minist. Agric. Cria, Invest. Pesq. Ser. Rec. Explot. Pesq. 1(5):175-204.

- SOKOLOVA, L.V. 1965. Distribution and biological characteristics of the main commercial fish of Campeche Bank. p. 208-224. In: A.S. Bogdanov (ed.). Sovetsko-Kubinskie rybokhozyaistvenne issledovaniya, VNIRO-CIP, Vypusk I. Isdatel 'stvo "Pishchevaya Promyshlennost", Moskva. Soviet-Cuban Fishery Research, Israel Program for Scientific Translations, Jerusalem, 1969.
- SSENTONGO, G.W. and P.A. LARKIN. 1973. Some simple methods of estimating mortality rates of exploited fish populations. J. Fish. Res. Board Can. 30:695-698.
- THRASH, B. 1985. Feasibility study of minced fish and surimi from Spanish sardine. NMFS S-K Grant Rep. no. 85-43, 7 p.
- UEBELHOER, M.R. 1984. Final report of results from the exploratory deepwater purse seining and refrigerated seawater chilling system demonstration project. Performed by: Raffield Fisheries, Inc., Port St. Joe, Fla., NMFS Award #NA82-GA-H-00007, 30 p.
- U. S. DEP. COMMER., NOAA. 1985. Gulf of Mexico coastal and ocean zones strategic assessment: data atlas. NOAA, Rockville, MD, unpag.
- WHITEHEAD, P.J.P. 1973. The clupeoid fishes of the Guianas. Bull. Brit. Mus. (Nat. Hist.) Zool. Suppl. 5:1-227.
- ZEI, M. 1969. The behavior of Sardinella aurita Val., in relation to light and temperature. In: A. Ben-Tuvia and W. Dickson (eds.). Proceedings of the FAO conference on fish behaviour in relation to fishing techniques and tactics. FAO Fish. Rep. 62, v. 3:469-475.

Table 1. Morphometric features of Sardinella aurita (N = 100) in the Gulf of Mexico (Prosvirov and Varea 1969).

| Character | Overall dimensions | M* | $\pm M^*$ | $\pm S^*$ |
|--|--------------------|-------|-----------|-----------|
| Body length (after Smith)*, (cm) | 16.5-19.7 | 17.95 | 0.064 | 0.64 |
| Weight, g | 55.1-99.7 | 73.12 | 0.875 | 8.75 |
| In % of body length (after Smith)* | | | | |
| Length of head | 22.3-25.0 | 23.43 | 0.055 | 0.55 |
| Maximal height of body | 18.5-24.5 | 22.60 | 0.114 | 1.14 |
| Minimal height of body | 6.0-8.0 | 6.78 | 0.041 | 0.41 |
| Antedorsal distance | 38.9-44.5 | 41.56 | 0.126 | 1.26 |
| Anteventral distance | 43.6-52.5 | 47.87 | 0.134 | 1.34 |
| Length of pectoral fin | 10.8-16.4 | 15.16 | 0.083 | 0.83 |
| Anteanal distance | 65.0-77.4 | 73.26 | 0.170 | 1.70 |
| Length of ventral fin | 7.1-9.4 | 8.36 | 0.042 | 0.42 |
| Distance between bases of anal and caudal fins | 23.2-29.6 | 26.46 | 0.122 | 1.22 |
| In % of length of the head | | | | |
| Length of snout | 27.3-39.6 | 33.64 | 0.226 | 2.26 |
| Diameter of eye | 20.0-25.7 | 23.50 | 0.114 | 1.14 |
| Postorbital section | 44.4-55.0 | 50.20 | 0.182 | 1.82 |
| Height of head | 62.9-79.0 | 72.57 | 0.340 | 3.40 |
| V | | | | |
| A | 16-18 | 17 | | |
| Number of rays B | 9 | 9 | | |
| D | 16-17 | | | |
| Number of vertebrae | 46 | 46 | | |

* Abbreviations are not defined in original paper, and Smith is not referenced, in Prosvirov and Varea (1969).

Table 2. Morphometric measurements expressed as percent of standard lengths for S. aurita (Hildebrand 1963).

| | |
|----------------------------------|--|
| Body depth: 21-26.9 | Pectoral fin: length 15-17.6 |
| Caudal peduncle: depth 7.7-10 | Distance from snout to origin of: |
| Head: length 22.3-27.2; | dorsal fin 40-43.5 |
| depth 17.6-19.7 | |
| Snout: length 6.0-7.5 | Scales: about 41-46; often lost from |
| Eye: diameter 5.3-7.3 | preserved specimens |
| Interorbital: width 4.2-5.3 | Ventral scutes: 32-34 |
| Maxillary: length 8.75-11 | Fin rays: dorsal 17-19, occasionally 20; |
| Anal fin: length of base 13.7-16 | anal 16-18; pectoral 15 or 16 |
| Pelvic fin: length 8.5-11 | Vertebrae: 45-57 (7 specimens) |

Table 3. List of features of larval S. aurita (Fahay 1983).

| | |
|---|-------------------|
| Spawning: September - February off Florida | Meristic features |
| Eggs - Pelagic, spherical | Myomeres: 45-48 |
| - Diameter: 0.94-1.40 mm | Vert: 16+29-31 |
| - Shell: smooth, transparent and thin | D : (15)16-19(20) |
| - Yolk: segmented | A : (14)16-17(19) |
| - Perivitelline space: moderate | Plv : 8-10 |
| - Oil globules: 1 | C : 8+10+9+7 |
| - O.G. diameter: 0.12-0.16 mm | |
| Larvae - Hatching occurs at about 3 mm | |
| - Body elongate with long straight gut; vent posterior to dorsal fin | |
| - Flexion occurs at about 11 mm TL, and transformation at about 25 mm TL | |
| - Last 2 anal rays become elongate in larger larvae | |
| - Larvae uncommon north of Cape Hatteras | |
| Important characters: | |
| - Myomeres between dorsal and anal fins 5-8; predorsal myomeres decrease from 28-24 | |
| - Dorsal pigment absent from notochord tip, but ventral pigment present | |
| - Anal fin ray count lower than in <u>Opisthonema</u> | |

Table 4. Summary of experimental midwater trawls (MWT) by U.S. research vessels in the northern Gulf of Mexico 1957-69.*

| Cruise code | Month year | Location | Gear** | Depth | Size of fish school & catch | Genus |
|-------------|---------------|---|-----------------------------|---------------------------------|---------------------------------|---|
| 01 42 | 1/57 | NE Gulf | 30 ft, 50 ft MWT | 20-90 fm | 300 yds + dia | Anchoa <u>Sardinella</u> <u>Decapterus</u> <u>Scomber</u> |
| 01 49 | 4/57 | Cape San Blas to Fort Myers | 40 ft MWT | 10-30 fm | 200 lbs/hr | Etrumeus <u>Anchoa</u> <u>Scomber</u> <u>Sardinella</u> |
| 01 57 | 2-3/59 | Mississippi River to Chandeleur Island Cedar Key to Cape Romano Cape San Blas | 40 ft MWT " " " | 8-12 fm 10-30 fm 10-12 fm | up to 1000 lbs/tow " " | Harengula <u>Opisthonema</u> <u>Anchoa</u> <u>Brevoortia</u> <u>Etrumeus</u> <u>Sardinella</u> |

* Information from Frederick Minkler, National Marine Fisheries Service, Pascagoula, MS, May 1985.

**Midwater trawl size in feet of headrope.

Table 5. Catch rate of S. aurita by depth and season*.

| Depth (fathoms) | December-February | | March-May | | June-August | | September-November | |
|--------------------|-------------------|--------|-------------------|------|-------------------|------|--------------------|------|
| | Number of tows | Hits** | Number of tows | Hits | Number of tows | Hits | Number of tows | Hits |
| 1-10 | 457 | 6 | 1175 | 19 | 2101 | 46 | 2251 | 44 |
| 11-20 | 647 | 16 | 1933 | 24 | 1156 | 39 | 2970 | 29 |
| 21-30 | 455 | 9 | 1065 | 7 | 615 | 16 | 2008 | 13 |
| 31-40 | 278 | 2 | 613 | 3 | 298 | 0 | 1421 | 2 |
| 41-50 | - | - | - | 1 | 206 | 1 | - | - |

* Data from Frederick Minkler, National Marine Fisheries Service, Pascagoula, MS, May 1985.

** Number of tows in which Spanish sardine was found.

Table 6. Catch rate of S. aurita by surface temperature (°C) and season*.

| Surface Temp. °C | December-February | | March-May | | June-August | | September-November | |
|---------------------|-------------------|--------|-------------------|------|-------------------|------|--------------------|------|
| | Number of tows | Hits** | Number of tows | Hits | Number of tows | Hits | Number of tows | Hits |
| 13 | 139 | 1 | 10 | 0 | - | - | 1 | 0 |
| 14 | 216 | 0 | 17 | 0 | - | - | 2 | 0 |
| 15 | 61 | 0 | 40 | 1 | - | - | - | - |
| 16 | 211 | 1 | 106 | 0 | - | - | 19 | 0 |
| 17 | 233 | 2 | 192 | 0 | 1 | 0 | 101 | 0 |
| 18 | 306 | 2 | 390 | 1 | - | - | 75 | 0 |
| 19 | 240 | 0 | 903 | 1 | 1 | 0 | 206 | 0 |
| 20 | 217 | 0 | 343 | 0 | - | - | 166 | 0 |
| 21 | 86 | 0 | 446 | 0 | 1 | 0 | 627 | 0 |
| 22 | 168 | 0 | 460 | 0 | 1 | 0 | 1088 | 0 |
| 23 | 134 | 0 | 298 | 6 | 17 | 0 | 1594 | 0 |
| 24 | 31 | 0 | 213 | 9 | 28 | 1 | 1244 | 0 |
| 25 | 1 | 0 | 48 | 2 | 26 | 1 | 384 | 1 |
| 26 | - | - | 77 | 2 | 279 | 0 | 699 | 8 |
| 27 | - | - | 109 | 1 | 549 | 6 | 538 | 15 |
| 28 | 3 | 0 | 25 | 1 | 1029 | 13 | 408 | 1 |
| 29 | 2 | 0 | 1 | 0 | 1009 | 77 | 392 | 0 |
| 30 | - | - | - | - | 367 | 1 | 64 | 1 |
| 31 | - | - | - | - | 236 | 4 | 21 | 0 |
| 32 | - | - | - | - | 41 | 0 | 1 | 0 |
| 33 | - | - | - | - | 4 | 0 | - | - |
| 34 | - | - | - | - | 1 | 0 | - | - |

* Data from Frederick Minkler, National Marine Fisheries Service, Pascagoula, MS, May 1985.

Shrimp and fish trawl (40 ft head rope) data from Gulf of Mexico 1950-1985.

** Number of tows in which Spanish sardine was found.

Table 7. Catch rate of S. aurita by bottom temperature (°C) and season*.

| Bottom Temp. °C | December-February | | March-May | | June-August | | September-November | |
|--------------------|-------------------|--------|-------------------|------|-------------------|------|--------------------|------|
| | Number of tows | Hits** | Number of tows | Hits | Number of tows | Hits | Number of tows | Hits |
| 17 | 45 | 0 | 129 | 1 | 42 | 0 | 35 | 0 |
| 18 | 84 | 0 | 249 | 0 | 75 | 0 | 58 | 0 |
| 19 | 64 | 0 | 294 | 0 | 147 | 0 | 78 | 0 |
| 20 | 19 | 0 | 148 | 0 | 123 | 0 | 58 | 0 |
| 21 | 35 | 0 | 149 | 0 | 281 | 0 | 145 | 0 |
| 22 | 54 | 0 | 53 | 0 | 238 | 1 | 303 | 1 |
| 23 | 87 | 0 | 59 | 0 | 219 | 8 | 448 | 0 |
| 24 | 93 | 0 | 18 | 0 | 164 | 5 | 517 | 0 |
| 25 | 5 | 0 | - | - | 86 | 0 | 131 | 0 |
| 26 | 9 | 0 | 3 | 0 | 266 | 3 | 163 | 2 |
| 27 | - | - | 2 | 0 | 164 | 1 | 82 | 2 |
| 28 | - | - | - | - | 118 | 0 | 15 | 1 |
| 29 | - | - | - | - | 60 | 2 | 3 | 0 |
| 30 | - | - | - | - | 9 | 0 | 2 | 0 |

* Data from Frederick Minkler, National Marine Fisheries Service, Pascagoula, MS, May 1985.

Shrimp and fish trawl (40 ft head rope) data from Gulf of Mexico 1950-1985.

** Percent of total number of tows in which Spanish sardine was present.

Table 8. Data on size, ovary weight, gonad index, fecundity and relative fecundity for Sardinella aurita (Martinez 1972).

| Fish | Standard length (cm) | Total length (cm) | Weight (g) | Ovary weight (g) | Gonad index (%) | Fecundity (no. ova) | Relative fecundity |
|------|----------------------------|-------------------------|---------------|------------------------|-----------------------|------------------------|-----------------------|
| 1 | 14.4 | 15.8 | 51.02 | 1.2100 | 2.3 | 59,682 | 1,198.0 |
| 2 | 14.4 | 15.6 | 50.17 | 2.2929 | 4.7 | 56,826 | 1,187.0 |
| 3 | 14.4 | 15.6 | 46.70 | 1.5288 | 3.3 | 61,815 | 1,370.0 |
| 4 | 14.6 | 16.1 | 49.93 | 1.7493 | 3.5 | 30,718 | 637.0 |
| 5 | 14.7 | 16.2 | 50.90 | .8847 | 1.7 | 45,824 | 916.0 |
| 6 | 15.0 | 16.2 | 53.81 | 2.1449 | 3.9 | 45,730 | 885.0 |
| 7 | 15.4 | 16.5 | 57.63 | 2.0681 | 3.5 | 86,269 | 1,552.0 |
| 8 | 15.5 | 16.8 | 63.91 | 2.7793 | 4.3 | 68,632 | 1,122.0 |

Table 9. Equations, standard deviations (S and S_b) and correlation coefficients (r) for the relations of fecundity (considering fecundity as the number of maturing and mature oocytes in the intermediate and most advanced mode) and relative fecundity to body weight and length in Spanish sardines, S. aurita (Martinez 1972).

| Variables | Results |
|---|--|
| Fecundity (F) and body weight (W) g | $F = 1,560 W - 25,764$; $S = 15,610$; $S_b = 1,086$; $r = .51$ |
| Fecundity (F) and standard length (SL) mm | $F = 18,939 SL - 223,367$; $S = 15,575$; $S_b = 13,088$; $r = .26$ |
| \log_{10} fecundity and standard length (SL) mm | $\log_{10} F = 2.789509 + 0.131643 SL$; $S = .13136$; $S_b = .1100388$; $r = .4382$ |
| \log_{10} fecundity and \log_{10} standard length (SL) mm | $\log_{10} F = 4.4593 \log_{10} SL - .4799$; $S = .13169$; $S_b = 3.7953$; $r = .43$ |
| Relative fecundity (F_r) and standard length (SL) mm | $F_r = 121.83 SL - 694.54$; $S = 307.45$; $S_b = 258.36$; $r = .19$ |

Table 10. Length, weight, batch fecundity, relative fecundity and gonad index of S. aurita from the Gulf of Mexico (Grall 1984).

| Pork length (mm) | Weight (g) | Batch fecundity (# of eggs) | Relative fecundity (eggs/g) | Gonad index (%) |
|------------------------|---------------|-----------------------------------|-----------------------------------|-----------------------|
| 146 | 45.9 | 21,214 | 505.1 | 8.50 |
| 154 | 57.0 | 31,737 | 623.5 | 10.70 |
| 162 | 60.5 | 26,721 | 476.3 | 7.27 |
| 165 | 67.7 | 45,099 | 733.3 | 9.16 |
| 166 | 74.5 | 34,462 | 500.2 | 7.52 |
| 168 | 67.7 | 67,433 | 1023.0 | 8.86 |
| 170 | 72.2 | 94,108 | 1484.4 | 12.19 |
| 175 | 88.4 | 69,550 | 899.7 | 12.56 |
| 177 | 73.8 | 36,842 | 545.8 | 8.54 |
| 179 | 80.6 | 92,405 | 1279.8 | 10.42 |
| 180 | 87.4 | 54,625 | 701.2 | 10.87 |
| 180 | 92.0 | 97,058 | 1202.2 | 12.17 |
| 182 | 87.1 | 80,816 | 1023.0 | 9.30 |
| 184 | 104.9 | 52,615 | 565.2 | 11.25 |
| 186 | 90.6 | 61,346 | 744.5 | 9.05 |
| 186 | 105.1 | 115,344 | 1227.1 | 10.55 |
| 187 | 107.6 | 98,449 | 1024.4 | 10.66 |
| 188 | 94.3 | 105,855 | 1232.3 | 8.91 |
| 188 | 107.5 | 146,728 | 1931.6 | 10.88 |
| 190 | 98.5 | 87,395 | 1232.3 | 9.24 |
| 193 | 98.7 | 40,022 | 443.7 | 8.61 |
| 195 | 114.5 | 41,541 | 399.0 | 9.08 |
| 195 | 118.9 | 47,802 | 437.0 | 7.99 |
| 198 | 120.9 | 139,800 | 1331.4 | 13.15 |
| 200 | 112.1 | 84,265 | 812.6 | 7.47 |
| average | | 70,312 | 871.7 | 9.80 |

Table 11. Length, weight, batch fecundity, relative fecundity and gonad index of S. aurita from the Atlantic Ocean (Grall 1984).

| Pork length (mm) | Weight (g) | Batch fecundity (# of eggs) | Relative fecundity (eggs/g) | Gonad index (%) |
|------------------------|---------------|-----------------------------------|-----------------------------------|-----------------------|
| 141 | 36.2 | 31,543 | 985.7 | 11.60 |
| 143 | 35.9 | 21,806 | 673.0 | 9.75 |
| 149 | 39.9 | 26,526 | 726.7 | 8.52 |
| 150 | 41.7 | 31,142 | 821.7 | 9.35 |
| 150 | 45.0 | 30,687 | 767.2 | 11.11 |
| 151 | 45.6 | 31,542 | 763.7 | 9.43 |
| 153 | 44.6 | 33,084 | 818.9 | 9.42 |
| 153 | 48.7 | 34,170 | 773.1 | 9.24 |
| 154 | 44.4 | 32,251 | 804.3 | 9.01 |
| 154 | 46.1 | 29,913 | 720.8 | 9.98 |
| 155 | 55.0 | 39,429 | 818.0 | 12.36 |
| 156 | 47.3 | 28,794 | 665.0 | 8.46 |
| 158 | 51.9 | 32,243 | 693.4 | 10.40 |
| 158 | 52.1 | 35,128 | 738.0 | 8.64 |
| 160 | 52.9 | 33,284 | 673.8 | 6.62 |
| 160 | 56.8 | 47,600 | 963.6 | 13.03 |
| 162 | 48.8 | 29,115 | 652.8 | 8.61 |
| 168 | 61.2 | 47,446 | 844.2 | 8.11 |
| 168 | 62.2 | 42,385 | 766.5 | 11.09 |
| 168 | 64.2 | 68,044 | 1195.8 | 11.37 |
| 168 | 65.5 | 40,822 | 684.9 | 9.01 |
| 176 | 77.6 | 41,186 | 580.1 | 8.51 |
| average | | 35,825 | 778.7 | 9.71 |

Table 12. Relationship of batch fecundity (F) and fork length (L in mm) and weight (W in g) for female S. aurita from the Gulf of Mexico (Grall 1984).

| Variables | Linear regression equation | Log-log linear regression equation |
|---|--|--|
| Batch fecundity vs fork length | $F = -165902.0 + 1317.5 L$ $r = 0.52$ $r^2 = 0.27$ $n = 25$ | $F = 350.4 \times 10^{-7} L^{4.1}$ $r = 0.62$ $r^2 = 0.39$ $n = 25$ |
| Batch fecundity vs fork length (adjusted) | $F = -318472.9 + 2227.1 L$ $r = 0.77$ $r^2 = 0.60$ $n = 20$ | $F = 118.8 \times 10^{-8} L^{6.1}$ $r = 0.83$ $r^2 = 0.68$ $n = 20$ |
| Batch fecundity vs weight | $F = -12389.5 + 934.74 W$ $r = 0.55$ $r^2 = 0.30$ $n = 25$ | $F = 158.4 W^{1.3}$ $r = 0.63$ $r^2 = 0.40$ $n = 25$ |
| Batch fecundity vs weight (adjusted) | $F = -46568.8 + 1413.7 W$ $r = 0.77$ $r^2 = 0.59$ $n = 20$ | $F = 20.99 W^{1.8}$ $r = 0.81$ $r^2 = 0.66$ $n = 20$ |

(*) Adjusted data include only females less than 190 mm FL.

Table 13. Relationship of batch fecundity (F), and fork length (L in mm) and weight (W in g) for female S. aurita from the Atlantic coast (Grall 1984).

| Variables | Linear regression equation | Log-log linear regression equation |
|--------------------------------|--|--|
| Batch fecundity vs fork length | $F = -80413.3 + 740.2 L$ $r = 0.67$ $r^2 = 0.45$ $n = 22$ | $F = 0.006 L^{3.1}$ $r = 0.71$ $r^2 = 0.51$ $n = 22$ |
| Batch fecundity vs weight | $F = -910.6 + 683.6 W$ $r = 0.73$ $r^2 = 0.53$ $n = 22$ | $F = 779.8 W^{0.97}$ $r = 0.79$ $r^2 = 0.63$ $n = 22$ |

Table 14. Mean fork length (FL) and weight (WT) of S. aurita in the Gulf of Mexico for 1981-1982 (Grall 1984).

| Year | Month | Mean FL (mm) | Mean WT (g) | Number of fish in sample |
|------|-------|--------------|-------------|--------------------------|
| 1981 | Apr | 166.77 | 64.89 | 289 |
| | May | 160.57 | 59.54 | 286 |
| | Jun | 170.00 | 69.44 | 882 |
| | Jul | 167.64 | 66.39 | 496 |
| | Aug | 157.56 | 57.95 | 310 |
| | Sep | 159.36 | 53.64 | 201 |
| | Oct | 123.54 | 26.41 | 590 |
| | Nov | 153.59 | 51.28 | 102 |
| 1982 | Apr | 134.63 | 29.13 | 109 |
| | May | 149.92 | 44.62 | 411 |
| | Jun | 162.60 | 60.71 | 820 |
| | Jul | 163.22 | 59.63 | 587 |
| | Aug | 164.26 | 62.58 | 204 |
| | Sep | 113.53 | 18.52 | 133 |
| | Oct | 120.78 | 21.86 | 371 |
| | Nov | 138.18 | 33.49 | 118 |

Table 15. Monthly mean fork length (FL) and weight (WT) of S. aurita catch taken by purse seine and by beach seine from the west coast of Florida for pooled years, 1981 and 1982 (Grall 1984).

| <u>Month</u> | <u>Purse seine</u> | | <u>Beach seine</u> | |
|--------------|--------------------|---------------|--------------------|---------------|
| | <u>FL (mm)</u> | <u>WT (g)</u> | <u>FL (mm)</u> | <u>WT (g)</u> |
| April | 144.21 | 42.00 * | 168.21 | 67.09 |
| May | 155.09 | 49.25 | 157.78 | 55.47 |
| June | 162.57 | 58.34 * | 168.19 | 67.65 |
| July | 161.74 | 56.89 * | 165.97 | 63.94 |
| August | 163.36 | 54.27 | 162.22 | 61.19 |
| September | - | - | 141.11 | 39.66 |
| October | 126.80 | 26.16 * | 120.93 | 24.59 |
| November | - | - | 148.23 | 44.80 |

* Significant difference in average length and weight between gear types at $P < 0.05$.

Table 16. Summary of the length-weight relationships (mm-g) for S. aurita on the Gulf coast of Florida (Grall 1984).

| <u>Year</u> | <u>Sex</u> | <u>Equation</u> | <u>r²</u> | <u>N</u> |
|-------------------|------------|--------------------------------------|----------------------|----------|
| 1981 | Males | $W = 12.80 \times 10^{-6} FL^{3.01}$ | 0.94 | 504 |
| | Females | $W = 5.55 \times 10^{-6} FL^{3.18}$ | 0.94 | 438 |
| | Combined | $W = 6.32 \times 10^{-6} FL^{3.15}$ | 0.96 | 3157 |
| 1982 | Males | $W = 1.07 \times 10^{-6} FL^{3.49}$ | 0.96 | 416 |
| | Females | $W = 0.87 \times 10^{-6} FL^{3.54}$ | 0.97 | 421 |
| | Combined | $W = 1.37 \times 10^{-6} FL^{3.45}$ | 0.96 | 2489 |
| 1981 + 1982 | Males | $W = 3.60 \times 10^{-6} FL^{3.26}$ | 0.95 | 917 |
| | Females | $W = 1.73 \times 10^{-6} FL^{3.40}$ | 0.96 | 884 |
| | Combined | $W = 3.45 \times 10^{-6} FL^{3.27}$ | 0.96 | 5913 |

Table 17. Summary of the length-weight relationships (mm-g) for S. aurita on the Atlantic coast of Florida (Grall 1984).

| Year | Sex | Equation | r^2 | N |
|------|----------|------------------------------------|-------|-----|
| 1981 | Males | $W=24.99 \times 10^{-6} FL^{2.85}$ | 0.83 | 189 |
| | Females | $W=3.08 \times 10^{-6} FL^{3.28}$ | 0.92 | 88 |
| | Combined | $W=3.82 \times 10^{-6} FL^{3.23}$ | 0.97 | 327 |

Table 18. Estimated age at length, back-calculated from the fork length-otolith regressions for S. aurita (Grall 1984).

| <u>Year</u> | <u>Area</u> | <u>Sex</u> | <u>Length (mm) at</u> | | |
|-------------|-------------|------------|-----------------------|--------------|--------------|
| | | | <u>Age 1</u> | <u>Age 2</u> | <u>Age 3</u> |
| 1981 | Gulf | Male | 104.26 | 147.41 | 165.85 |
| | | Female | 103.91 | 153.94 | 171.09 |
| | | Combined | 104.01 | 150.86 | 169.84 |
| 1982 | Gulf | Male | 107.26 | 146.14 | 170.34 |
| | | Female | 110.74 | 149.38 | 172.10 |
| | | Combined | 110.36 | 147.01 | 170.12 |
| 1981 | Gulf | Male | 104.98 | 146.86 | 168.32 |
| + | | Female | 105.96 | 151.71 | 171.11 |
| 1982 | | Combined | 105.19 | 149.32 | 170.12 |
| 1981 | Atlantic | Male | 111.97 | 141.87 | --- |
| | | Female | 111.68 | 144.44 | --- |
| | | Combined | 110.20 | 142.75 | --- |

Table 19. Distribution of back-calculated length (mm) at annulus (BC) and length (mm) at capture (AC) for S. aurita in the Gulf of Mexico, males and females combined in pooled years, 1981 and 1982 (Grall 1984).

| Length class (mm) | <u>Annulus 1</u> | | <u>Annulus 2</u> | | <u>Annulus 3</u> | |
|-------------------|------------------|-----------|------------------|-----------|------------------|-----------|
| | <u>BC</u> | <u>AC</u> | <u>BC</u> | <u>AC</u> | <u>BC</u> | <u>AC</u> |
| 70 - 79 | 10 | | | | | |
| 80 - 89 | 46 | | | | | |
| 90 - 99 | 80 | 1 | | | | |
| 100 - 109 | 107 | 8 | | | | |
| 110 - 119 | 86 | 45 | 5 | | | |
| 120 - 129 | 90 | 53 | 8 | | | |
| 130 - 139 | 39 | 73 | 39 | 1 | | |
| 140 - 149 | 23 | 100 | 125 | 18 | 3 | |
| 150 - 159 | 10 | 182 | 88 | 87 | 7 | 1 |
| 160 - 169 | | 191 | 59 | 187 | 19 | 1 |
| 170 - 179 | | 51 | 19 | 143 | 24 | 25 |
| 180 - 189 | | 11 | 1 | 64 | 2 | 37 |
| 190 - 199 | | 1 | | 6 | | 12 |
| 200 - 209 | | 1 | 1 | 1 | | |
| <u>N</u> | 491 | 717 | 343 | 507 | 55 | 76 |
| <u>x</u> | 105.19 | 150.39 | 149.32 | 167.55 | 170.12 | 181.67 |
| <u>st. dev.</u> | 16.8 | 17.3 | 11.7 | 10.2 | 8.2 | 7.2 |

Table 20, Back-calculated length at age for S. aurita based on the premise that the annulus is formed at six months rather than one year (Grall 1984).

| <u>Year</u> | <u>Area</u> | <u>Sex</u> | <u>Length (mm) at</u> | | |
|-------------|-------------|------------|-----------------------|--------------|--------------|
| | | | <u>Age 1</u> | <u>Age 2</u> | <u>Age 3</u> |
| 1981 | Gulf | Male | 129.49 | 159.49 | 174.19 |
| | | Female | 133.27 | 165.52 | 180.11 |
| | | Combined | 131.68 | 163.43 | 178.55 |
| 1982 | Gulf | Male | 131.22 | 161.91 | 179.18 |
| | | Female | 127.61 | 160.26 | 181.42 |
| | | Combined | 129.11 | 160.99 | 179.93 |
| 1981 | Gulf | Male | 129.28 | 159.43 | 174.81 |
| + | | Female | 132.82 | 164.49 | 179.68 |
| 1982 | | Combined | 130.75 | 162.65 | 179.20 |

Table 21. Monthly age (in years) distribution by number of S. aurita for males (M) and females (F) (from Grall 1984).

| <u>Year</u> | <u>Month</u> | <u>Age 0.0</u> | | <u>Age 0.5-1.0</u> | | <u>Age 1.5-2.0</u> | | <u>Age 2.5-3.0</u> | |
|-------------|--------------|----------------|----------|--------------------|-----------|--------------------|----------|--------------------|----------|
| | | <u>M</u> | <u>F</u> | <u>M</u> | <u>F</u> | <u>M</u> | <u>F</u> | <u>M</u> | <u>F</u> |
| 1981 | Apr | 13 | 7 | 17 | 4 | 6 | 13 | 1 | 3 |
| | May | 18 | 12 | 29 | 28 | 14 | 9 | 3 | 9 |
| | Jun | 10 | 4 | 71 | 49 | 53 | 62 | 5 | 9 |
| | Jul | 0 | 2 | 35 | 41 | 20 | 55 | 2 | 6 |
| | Aug | 10 | 1 | 41 | 8 | 14 | 9 | 0 | 2 |
| | Sep | 8 | 1 | 25 | 11 | 8 | 3 | 0 | 0 |
| | Oct | 70 | 35 | 13 | 9 | 0 | 1 | 0 | 0 |
| | Nov | <u>9</u> | <u>6</u> | <u>5</u> | <u>2</u> | <u>3</u> | <u>3</u> | <u>0</u> | <u>0</u> |
| Total | | 138 | 68 | 236 | 152 | 118 | 155 | 11 | 29 |
| | | | | | | | | | |
| 1982 | Apr | 12 | 9 | 7 | 2 | 0 | 0 | 0 | 0 |
| | May | 3 | 4 | 42 | 30 | 12 | 8 | 0 | 0 |
| | Jun | 3 | 1 | 46 | 31 | 46 | 57 | 16 | 13 |
| | Jul | 3 | 2 | 32 | 34 | 38 | 29 | 1 | 1 |
| | Aug | 1 | 1 | 10 | 7 | 20 | 11 | 1 | 0 |
| | Sep | 19 | 22 | 7 | 7 | 0 | 0 | 0 | 0 |
| | Oct | 6 | 11 | 19 | 29 | 0 | 0 | 0 | 0 |
| | Nov | <u>2</u> | <u>1</u> | <u>10</u> | <u>14</u> | <u>2</u> | <u>3</u> | <u>0</u> | <u>0</u> |
| Total | | 49 | 51 | 173 | 154 | 118 | 110 | 18 | 14 |

Table 22. Length and estimated age at capture of S. aurita determined from the von Bertalanffy growth model for pooled years, 1981-1982 (Grall 1984).

| <u>Sex</u> | <u>Mean length at capture (mm)</u> | <u>Age (yrs)</u> | <u>Adjusted age*</u> | <u>Frequency</u> |
|-------------|--|------------------|----------------------|------------------|
| Males | 132.6 | 1.63 | 1.08 | 187 |
| | 153.7 | 2.27 | 1.75 | 409 |
| | 164.8 | 2.37 | 2.28 | 236 |
| | 179.0 | 3.33 | 3.45 | 29 |
| <u>Mean</u> | | <u>2.35</u> | <u>2.01</u> | |
| Females | 132.2 | 1.57 | 0.99 | 119 |
| | 154.0 | 2.03 | 1.58 | 306 |
| | 170.0 | 2.27 | 2.28 | 265 |
| | 183.1 | 3.23 | 3.38 | 43 |
| <u>Mean</u> | | <u>2.22</u> | <u>2.10</u> | |
| Combined | 127.5 | 1.60 | 0.93 | 378 |
| | 151.2 | 2.10 | 1.56 | 779 |
| | 167.5 | 2.30 | 2.23 | 502 |
| | 181.6 | 3.30 | 3.22 | 73 |
| <u>Mean</u> | | <u>2.24</u> | <u>1.90</u> | |

* The adjusted age is calculated from the von Bertalanffy curve which is based on the premise that the first annulus is formed at six months rather than one year.

Table 23. Growth parameters of S. aurita from different areas (Grall 1984).

| Author | Area | L_{∞} (mm) | K | t_0 |
|----------------------------------|-----------|-------------------|--------|--------|
| Boely, 1982 | Senegal | 306.3 FL | 1.2060 | -.0620 |
| El-Maghraby <u>et al.</u> , 1970 | Egypt | 260.0 FL | 0.5310 | .3395 |
| Gheno, 1975 | Congo | 259.7 FL | 1.0230 | .0885 |
| Gheno, 1975 | Congo | 260.0 FL | 1.2090 | -.0296 |
| Heald, 1966 | Venezuela | 241.5 SL | 0.3200 | ----- |
| Krzeptowski, 1981 | Senegal | 331.5 FL | 0.2832 | -.9890 |
| Matsuura, 1977 | Brazil | 244.0 FL | 0.4414 | ----- |
| Grall (1984) | Florida | 220.1 FL | 0.4708 | -.3810 |
| | | 197.1 FL | 0.6560 | -.6604 |

Table 24. Mortality rate estimates (M) and annual survival (S) of S. aurita based on the methods of Pauly (1980) and Ssentongo and Larkin (1973) for the years 1981-1982 (Grall 1984).

| Method | Males | Females | Combined |
|--------------------------|------------|------------|------------|
| Pauly | M = 0.5995 | M = 0.5608 | M = 0.5769 |
| | S = 0.5491 | S = 0.5707 | S = 0.5616 |
| Pauly (*) | M = 0.7588 | M = 0.8001 | M = 0.7392 |
| | S = 0.4682 | S = 0.4493 | S = 0.4775 |
| Ssentongo & Larkin | M = 0.8916 | M = 1.009 | M = 0.9894 |
| | S = 0.4100 | S = 0.3646 | S = 0.3718 |
| Ssentongo & Larkin(*) | M = 0.7988 | M = 0.7709 | M = 0.8433 |
| | S = 0.4499 | S = 0.4694 | S = 0.4303 |

(*) Represents mortality and survival estimates based on the adjusted growth parameters.

Table 25. Yield-per-recruit, mean weight, length and age of S. aurita in the catch at A.) maximum fishing mortality, B.) optimum fishing mortality and C.) maximum exploitation rates (Grall 1984).

| Data used | F | E | Yield per recruit (g) | Mean weight (g) | Mean length (mm FL) | Mean age (yr) |
|-----------|------|------|-----------------------------|-----------------------|---------------------------|---------------------|
| A.) | | | | | | |
| original | 6.7 | 0.87 | 24.14 | 27.70 | 129.28 | 1.49 |
| adjusted | 3.8 | 0.82 | 24.42 | 29.84 | 132.25 | 1.03 |
| B.) | | | | | | |
| original | 1.5 | 0.60 | 21.91 | 36.52 | 140.80 | 1.58 |
| adjusted | 1.5 | 0.64 | 23.15 | 36.16 | 140.38 | 1.24 |
| C.) | | | | | | |
| original | 1.0 | 0.50 | 19.87 | 39.58 | 145.60 | 1.92 |
| adjusted | 0.85 | 0.50 | 20.39 | 40.78 | 145.63 | 1.39 |

Table 26. Some predators of S. aurita.

| Predator | Location studied | Spanish sardine occurrence in stomachs | | Reference |
|---------------------|---------------------------|---|-------------------|-----------------------------|
| | | Percent frequency | Percent volume | |
| Bluefish | Carolinas to Louisiana | 0 - 2.7 | 0 - 2.9 | Naughton & Saloman 1984 |
| Spanish mackerel | Carolinas to Texas | 0.5 - 3.2 | 2.2 - 20.4 | Saloman & Naughton 1983b |
| Crevalle jack | Florida to Texas | 0 - 6.2 | 0 - 9.3 | Saloman & Naughton 1984 |
| King mackerel | Carolinas to Texas | 0 - 9.0 | 0 - 24.3 | Saloman & Naughton 1983a |

Table 27. Species composition of purse seine catches, Florida, October, 1981 (Roithmayr 1983).

| Species | Tons | Percent composition |
|--------------------------------|------|---------------------|
| Thread herring | 121 | 41 |
| Ladyfish | 57 | 19 |
| Spot | 48 | 16 |
| Menhaden | 46 | 15 |
| Spanish sardine and round scad | 27 | 9 |
| TOTAL | 299 | |

Table 28. "Typical" drum seine conversion cost (projected) (Harrell 1982).

| | |
|--|-----------|
| Deck Equipment | \$ 27,500 |
| Drum Seine | 9,400 |
| Spool Assembly | 8,000 |
| Miscellaneous Items | 7,500 |
| Net | 50,000 |
| Total Cost | \$102,400 |
| (3rd Quarter 1981 dollars are estimated) | |

Table 29. Projected drum seine operating costs (Harrell 1982).

Assumptions (Basis):

1. Diesel Fuel Cost (Anderson Results): \$1.18/gallon
2. Fuel Consumption (Clark Results): 0.023 gallons per pound
3. Life of Drum Seine Mechanical Equipment: 12 years
4. Life of Electronic: 7 years (Due principally to obsolescence)
5. Life of Net: 2 years
6. Original Cost of Mechanical Equipment: \$50,000
7. Original Cost of Electronics: \$30,000
8. Estimated Insurance Cost: \$10,000
9. Ice Consumption: 0.765 pounds of ice per pound of fish (Clark actual results)
10. Ice Cost: \$4.75 per block (300 pounds)
11. Labor & Spotter Plane Cost: Divide Total Gross Stock as follows:
 48.6% - Vessel 10% - Spotter Plane
 32.4% - Crew 9% - Captain
12. Original Cost of Seine: \$50,000
13. Estimated Maintenance Cost: \$12 - 15,000 per year per boat (Use 2 boats)
14. Estimate Harvesting Rate: 33,000 pounds per day (Assume Clark Seafood's rate is typical)

Using these assumptions, the following operating costs can be expected
 (Note: per pound basis assumes boat produces 3,000,000 pounds of fish per year with 23 cents per pounds).

| <u>Cost Component</u> | <u>Annual Cost (Dollars)</u> | <u>(Cost per Pound (Cents))</u> |
|----------------------------|------------------------------|---------------------------------|
| Vessel depreciation | 15,000 | 0.500 |
| Mechanical depreciation | 4,167 | 0.139 |
| Electronic depreciation | 4,285 | 0.142 |
| Line depreciation | 25,000 | 0.833 |
| Vessel cost | 69,000 | 2.300 |
| cost | 47,500 | 1.583 |
| Insurance | 10,000 | 0.333 |
| Maintenance | 15,000 | 0.500 |
| Harbor & spotter plane | 292,560 | 9.752 |
| (Percent of gross revenue) | | |
| Miscellaneous | <u>5,000</u> | <u>0.167</u> |
| TOTAL COST | \$487,512 | 16.25 cents/lb |

Table 30. Sample comparison of drum seine and shrimp trawl costs and earnings (Harrell 1982).

| <u>Cost component</u> | <u>Drum seine</u> | <u>Shrimp trawl</u> |
|--|-------------------|---------------------|
| 1. Vessel Characteristics: | | |
| Gross Tonnage | 80 | 108 |
| Length | 70 | 69 |
| Horsepower | 245 | 352 |
| Crew Size | 7 | 3 |
| 2. Gross Revenue | \$450,000 | 150,000 |
| 3. Variable Costs: | | |
| (a) Goods & Services | 134,500 | 65,700 |
| (b) Labor | 145,800 | 50,000 |
| (c) Opportunity Cost of Capital | 2,018 | 985 |
| Total Variable Costs | 282,318 | 116,685 |
| 4. Fixed Costs: | | |
| (a) Insurance | 10,000 | 7,000 |
| (b) Miscellaneous | 5,000 | 2,086 |
| (c) Depreciation (See Table #8) | 48,452 | 8,667 |
| Total Fixed Costs | 63,452 | 17,753 |
| 5. Total Costs (3 ÷ 4) | 345,770 | 134,438 |
| 6. Returns to operator's labor, management and capital (2-5) | 104,230 | 15,562 |
| 7. Opportunity Cost of Management | 36,000 | 15,000 |
| 8. Returns to Capital (6-7) | 68,230 | 562 |
| 9. Total Investment | 325,000 | 275,000 |
| 10. Return on Investment (%) | 21.0 | 0.2 |
| Drum seine assumptions: | | |
| Annual Catch: 3,000,000 pounds | | |
| Catch Value: 15 cents per pound | | |

Table 31. Species composition, weight, and value of purse-seined fish in 1982 (Uebelhoer 1984).

| Species | No. of sets | Amount harvested (lbs) | Unit value | Total value |
|-----------------|-------------|------------------------|-----------------------------------|--------------------|
| Spanish Sardine | 4 | 217,678 (15.9%) | 6¢/lb. | \$13,060.68(11.0%) |
| Ladyfish | 5 | 425,050 (31.0%) | 8¢/lb. | 34,004.00(30.0%) |
| Mullet | 1 | 7,100 (0.5%) | 15¢/lb. | 1,065.00(0.9%) |
| Bonito | 3 | 172,706 (13.9%) | 8¢/lb. | 13,816.48(12.2%) |
| Jack Crevalle | 6 | 198,440 (14.5%) | 8¢/lb. | 15,875.20(14.0%) |
| Menhaden | 1 | 140,000 (10.2%) | 4.25¢/lb. | 5,950.00(5.3%) |
| Blue Runner | 3 | 93,500 (6.8%) | 17.0¢/lb. | 15,895.00(14.0%) |
| Red Fish | 2 | 72,000 (5.3%) | 12.0¢/lb. | 8,640.00(7.6%) |
| Blue Fish | 1 | 26,000 (1.9%) | 18.0¢/lb. | 4,680.00(4.1%) |
| Total | 26 | 1,352,474 | 8.25¢/lb (Weighted Average) | \$113,066.36 |

Table 32. Production and latent fishery resource potentials of the Gulf of Mexico in 1968
(Bullis and Carpenter 1968).

| | Production (Millions of Pounds) | Latent (Millions of Pounds) | Expansion factor | Species of greatest potential |
|-------------------------|---------------------------------------|-----------------------------------|---------------------|--|
| Coastal Pelagic Fish | 1,064 | 8,448 | 8 | Herring Sardine Anchovy |
| Bottomfish (industrial) | 88 | 5,673 | 64 | Croaker Spot Seatrout |
| Bottomfish (food) | 54 | 1,005 | 19 | Snapper Grouper |
| Shellfish | 257 | 2,835 | 11 | Scallops Squid Lobsters Crabs Shrimp |
| Mid-water Fish | None | 2,148 | 100+ | Butterfish Kumper Scad Harvest fish |
| High Seas Pelagic Fish | None | 912 | 100+ | Sharks Tuna Flying fish Dolphin |
| Total | 1,463 | 21,021 | 14 | |

Table 33. Status of the coastal herrings and associated stocks in the Gulf of Mexico and south Atlantic (Reintjes 1979).

| <u>Species</u> | <u>Total estimated catch</u> | <u>Biomass</u> | <u>Potential Yield</u> | <u>Available for expansion</u> |
|--------------------------------------|--------------------------------------|----------------|----------------------------|--|
| -----thousand metric tons----- | | | | |
| Atlantic thread herring ^d | 25-50 | 110-370 | 60-120 | 35-70 |
| Spanish sardine ^d | .5 | 110-370 | 60-120 | 60-120 |
| Scaled sardine ^d | .5 | 185 | 46-92 | 46-92 |
| Round herring | .25 | 130-700 | 50-250 | 50-250 |
| Bay anchovy | b | a | a | a |
| Striped anchovy | b | a | a | a |
| Silver anchovy | b | a | a | a |
| Round scad ^d | 1. | 100-200 | 70-85 | 69-84 |
| Rough scad | 0 | 20-50 | 12-14 | 12-14 |
| Atlantic bumper | 2.5 | a | a | a |
| Total ^c | 30-55 | 655-1,875 | 298-681 | 272-630 |

a Unknown

b Shrimp discards represent about 17,000 tons of anchovies.

c Numbers rounded

d Directed fisheries primarily limited to Florida

Table 34. Composition of *S. aurita*.

| Reference | Description | Moisture % | Protein % | Fat % | Ash % | Carbo- hydrates % | Calories Per 100 gm | Phosphorus mg/100 g | Chlorine mg | Arsenic parts per million |
|------------------------------|---|--|-------------------------|------------------------|----------------------|-------------------------|------------------------|------------------------|----------------|---------------------------------|
| Sidwell 1981 | Muscle Raw | \bar{x} 71.0 Range 67.9-74.8 n 4 | 22.9 18.3-32.6 5 | 6.0 2.3-9.9 6 | 1.7 1.1-2.1 5 | 0 | 146 | 331 | 140 | 0.99 |
| | Whole Raw | 71.6 | 19.0 18.3-19.6 2 | 3.6 | 6.5 | 0 | 108 | | | |
| | Dried | 23.9 22.8-24.9 2 | 57.0 | 5.7 | 14.5 | 0 | 279 | | | |
| | Salted & smoked | 23.3 22.5-24.1 2 | 56.3 | 5.4 | 14.2 | 0.8 | 277 | | | |
| Herzberg and Pasteur 1969 | Edible parts with skeletal frame | | 20.7 17.0-22.3 | 3.7 0.4-20.0 | 1.9 1.4-2.9 | | | | | |
| Sidwell et. al 1974 | Not listed | 74.8 65.9-79.9 50 | 20.5 17.3-22.3 49 | 3.8 0.4-20.0 125 | 2.1 1.4-2.9 49 | | | | | |

Table 35. Major fatty acids of raw and canned S. aurita (weight percent composition) (Hale and Brown 1983).

| Fatty acid | Sample form | | |
|--------------------------|-------------|--|----------------------------|
| | Raw | Canned ($F_0 = 13.3$) ^{1/} | Canned ($F_0 = 17.7$) |
| 14:0 | 3.9 | 3.5 | 4.1 |
| 16:0 | 22.2 | 21.8 | 22.8 |
| 17:0 | 1.6 | 1.5 | 1.5 |
| 18:0 | 7.5 | 7.7 | 7.3 |
| 16:1 | 5.7 | 5.4 | 6.1 |
| 18:1 | 11.3 | 10.2 | 11.4 |
| 20:1 | 1.3 | 1.2 | 1.1 |
| 22:1 | 0.1 | -- | 0.1 |
| 18:2 ω 6 | 1.2 | 1.3 | 1.4 |
| 18:3 ω 3 | 0.7 | 0.8 | 0.8 |
| 18:4 ω 3 | 0.7 | 0.6 | 0.8 |
| 20:4 ω 6 | 1.8 | 1.9 | 1.8 |
| 20:5 ω 3 | 8.1 | 7.8 | 9.4 |
| 22:5 ω 6 | 1.2 | 1.4 | 1.2 |
| 22:5 ω 3 | 1.4 | 1.2 | 1.3 |
| 22:6 ω 3 | 22.7 | 25.1 | 22.1 |
| Total saturated | 37.7 | 37.3 | 39.0 |
| Total monogenes | 19.4 | 17.8 | 19.5 |
| Total PUFA ^{2/} | 42.9 | 44.9 | 41.6 |
| Total HUFA ^{3/} | 33.6 | 35.6 | 33.9 |

^{1/} F_0 = measure of total heat exposure at the center of the can.

^{2/} PUFA = polyunsaturated fatty acids.

^{3/} HUFA = highly unsaturated fatty acids.

Table 36. Effect of packing media on concentrations of total lipids and selected fatty acids for canned S. aurita (Hale and Brown 1983).

| Item | Sample | | |
|-----------------|--------------------------|--------------------|-----------------------|
| | Raw, headed and gutted | Canned in 2% brine | Canned in soybean oil |
| | | Percent | |
| Total lipids | 1.3 | 2.05 | 9.64 |
| | Fatty acids (% of total) | | |
| 16:0 | 25.18 | 25.38 | 13.81 |
| 18:0 | 8.09 | 8.48 | 5.04 |
| 18:1 | 9.71 | 11.74 | 21.42 |
| 18:2 ω 6 | 1.06 | 1.32 | 43.50 |
| 20:5 ω 3 | 4.86 | 4.29 | 0.84 |
| 22:6 ω 3 | 29.46 | 25.50 | 4.34 |

Table 37. Lipid class composition of raw and canned S. aurita (Hale and Brown 1983).

| Lipid class | Sample form | | |
|------------------|------------------|--------------------------------|--------------------------------|
| | Raw | Canned (F _O = 13.3) | Canned (F _O = 17.7) |
| | Percent of total | | |
| Triglycerides | 67 | 55 | 66 |
| Free fatty acids | 7 | 7 | 5 |
| Sterols | 2 | 3 | 2 |
| Polar lipids | 22 | 35 | 26 |
| Unidentified | 2 | -- | -- |

Table 38. Size data for whole S. aurita, and proximate chemical compositions for dressed, headed and gutted (H & G) fish, both raw and canned (Hale and Brown 1983).

| | Size data (whole fish) | |
|---------------------|---|---------------|
| Item | <u>Spanish sardine</u> | |
| Mean weight (g±SD) | 71.5 ± 23.8 | |
| Mean length (cm±SD) | 16.9 ± 1.7 | |
| | <u>Proximate composition (dressed, H&G)</u> | |
| | <u>Raw</u> | <u>Canned</u> |
| Moisture (%) | 75.41 | 69.63 |
| Protein (%) | 20.09 | 24.52 |
| Lipid (%) | 3.22 | 4.21 |
| Ash (%) | 1.75 | 1.90 |

Table 39. Per capita consumption of sardines (Perkins 1981).

| | Pounds Per capita |
|------|-------------------|
| 1970 | .4 |
| 1971 | .4 |
| 1972 | .4 |
| 1973 | .5 |
| 1974 | .4 |
| 1975 | .2 |
| 1976 | .3 |
| 1977 | .3 |
| 1978 | .3 |
| 1979 | .3 |

Table 40. Hypothetical price and quantity demanded relationships with increased domestic production of canned sardines (Perkins 1981).

| <u>Quantity</u> | <u>Price</u> | <u>Total revenue</u> |
|-----------------|--------------|----------------------|
| 000 lbs. | Dol. | 000 dol. |
| 30,000 | \$1.50 | \$45,000 |
| 30,300 | 1.49 | 45,147 |
| 30,600 | 1.48 | 45,288 |
| 30,900 | 1.47 | 45,423 |
| 31,200 | 1.46 | 45,552 |

Table 41. Hypothetical per capita consumption of domestic sardines (Perkins 1981).

| <u>Production</u> | <u>Price</u> | <u>Consumption of sardines</u> ¹ |
|-------------------|--------------|---|
| Mil. lbs. | Dol. per lb. | lbs. |
| 30,000 | \$1.50 | .11 |
| 30,300 | 1.49 | |
| 30,600 | 1.48 | |
| 30,900 | 1.47 | |
| 31,200 | 1.46 | |
| 60,000 | .91 | .22 |

¹ Assuming unchanged consumption of imported canned sardines.

Table 42. Gulf of Mexico and south Atlantic export firms offering specified under-utilized species for export (Centaur Associates 1981).

| Product for export | Export firms | |
|-------------------------|--------------|---------------------------|
| | Number | Percent of firms surveyed |
| Black mullet | 27 | 79 |
| Bluefish | 21 | 62 |
| Jack crevalle | 20 | 59 |
| Little tunny | 19 | 56 |
| Bluerunner | 18 | 53 |
| Ladyfish | 18 | 53 |
| Spanish sardine | 17 | 50 |
| Sheepshead | 15 | 44 |
| Rock shrimp | 15 | 44 |
| Black drum | 14 | 41 |
| Atlantic thread herring | 12 | 35 |
| Sea catfish | 10 | 29 |
| Red porgy | 9 | 26 |
| Atlantic croaker | 9 | 26 |
| Blackfin tuna | 7 | 21 |
| Spot | 7 | 21 |
| Round herring | 7 | 21 |
| Black mullet roe | 6 | 18 |
| Gulf quahog | 2 | 6 |

Table 43. Locally produced seafood products exported from the southeast, 1980. (Prochaska and Cato 1981).

| Seafood product | Total | Percent of exporters reporting sales | |
|-----------------|-------|--------------------------------------|-------------|
| | | Large firms | Small firms |
| Mullet | 62 | 100 | 29 |
| Mullet Roe | 46 | 67 | 29 |
| Shrimp | 31 | 33 | 29 |
| Mackerel | 23 | 50 | 0 |
| Red Drum | 23 | 50 | 0 |
| Scallops | 8 | 17 | 0 |
| Monkfish | 8 | 17 | 0 |
| Squid | 8 | 17 | 0 |
| Butterfish | 8 | 17 | 0 |
| Ladyfish | 15 | 33 | 0 |
| Jack crevalle | 15 | 33 | 0 |
| Sardines | 8 | 0 | 14 |
| Porgy | 8 | 0 | 14 |
| King Mackerel | 15 | 33 | 0 |
| Lobster | 15 | 33 | 0 |
| Blue fish | 8 | 17 | 0 |
| Oyster stew | 8 | 0 | 14 |
| Black drum | 8 | 17 | 0 |
| Sheepshead | 8 | 17 | 0 |

^a Firm size based on total volume of sales and not sales of individual species. Small firms are those with sales less than one million pounds.






Figure 1. Diagram of a Spanish sardine, Sardinella aurita (Fischer 1978).



S. brasiliensis *S. aurita*
LOWER LIMBS OF GILL ARCHES



Figure 2. Identification characters for *S. aurita* (Fischer 1978).




Figure 3. Larval S. aurita as illustrated by Fahay (1983).




Figure 4. Larval *S. aurita* as illustrated by Jones et al. (1978).




Figure 5. Distribution of *S. aurita* in the Gulf of Mexico and adjacent areas (U.S. Dep. Commer., NOAA 1985).




Figure 6. Stations (solid circles) at which *Sardinella aurita* (= *S. anchovia*) larvae occurred at least once during 17 cruises to the eastern Gulf of Mexico, 1971-1974 (Houde et al. 1979).




Figure 7. Distribution and abundance of *S. aurita* larvae in the eastern Gulf of Mexico, 1971-1974. 0 = tows in which larvae occurred (Houde et al. 1979).




Figure 7. (Continued).




Figure 8. Estimated mean abundance (number under 10 m² of sea surface) of *S. aurita* larvae (Houde et al. 1979).




Figure 9. Catch rates for exploratory fishing for *S. aurita*, 1950 to 1985. a) Fall-winter (0=fall; 1=winter) b) spring-summer (0=spring; 1=summer). Also, 0 and 1 are bottom trawls; 4 are mid-water trawls. Numbers are pounds/hr. Information from Frederick Minkler, NMFS, Pascagoula Laboratory, Miss., May 1985.




Figure 10. Regression of fecundity, F , on weight for *S. aurita* (Martinez 1972).




Figure 11. The relationship between the gonad index and the stage of maturity measured in 400 females of *S. aurita*. Maturity stage (6) was not plotted due to insufficient data (Grall 1984).




Figure 12. Graphic relationship between fecundity and fork length of 26 female *S. aurita* from the west coast of Florida. The dashed line (A) was fitted to all data points including females which were longer than 190 mm (indicated by solid dots). The solid line (B) was fitted to the open dots which represented females less than 190 mm (sic) in length (Grall 1984).




Figure 13. Graphic relationship between fecundity and weight of 26 female *S. aurita* from the west coast of Florida. The dashed line (A) was fitted to all data points including females which were longer than 190 mm (indicated by solid dots). The solid line (B) was fitted to the open dots which represented females less than 190 mm in length (Grall 1984).





Figure 14. Frequency of mature *S. aurita* by fork length (Graff 1984).




Figure 15. Mean gonad index by month for male and female *S. aurita* during 1981-1982 (Graill 1984).






Figure 16. Annual length frequency of *S. aurita* for years 1980-1982. Distributions (A), (B) and (C) represent samples from the west coast of Florida and (D) represents samples from the east coast of Florida (Grall 1984).



17a. Monthly length frequencies of *S. aurita* collected from the west coast of Florida during 1981 (Grall 1984).



17b. Monthly length frequencies of *S. aurita* collected from the west coast of Florida during 1982 (Grall 1984).




Figure 18. Mean fork length (mm) and weight (g) by month of S. aurita (Grall 1984).




Figure 19. Length-weight relationships of *S. aurita* from the east and west coasts of Florida (Graff 1984).




Figure 20. The von Bertalanffy growth curve fitted to age data based on first annulus formation at approximately (A) one year of age or (B) six months of age of *S. aurita* (Grall 1984).




Figure 21. Yield-per-recruit curves assuming several ages at first capture (t) under various fishing mortalities (F) based on approximately first annulus formation at (A) one year of age or (B) six months of age of *S. aurita* (Grall 1984).




Figure 22. Apparent abundance by length of *S. aurita* larvae, collected on ichthyoplankton survey cruises to the eastern Gulf of Mexico, 1971, 1972, and 1973 (Houde et al. 1979).




Figure 23. The use of gill nets for catching herring: A. standard surface floating gill net set at night; B. use of gill net to encircle herring during the day; C. use of lights to attract herring to net (Harvey 1982).




Figure 24. Beach seine in operation: A. plane view; B. lateral view (Harvey 1982).




Figure 25. Sequence of purse seine operation: 1) setting the net 2) pursing 3) haul in purse seine rings and leadline (Roithmayr 1983).




Figure 26. Commercial purse seine grounds (Roithmayr 1983) .




Figure 27. Price and per capita consumption of sardines, 1970-1979 (Perkins 1981).




Figure 28. Demand curve for sardines (Perkins 1981).




Figure 29. Summary of information needs concerning coastal herrings (Christmas et al. 1985).